



Review Article

Studies of Grafts in vegetables, an alternative for agricultural production under stress conditions: Physiological responses

Jose Bernal Alzate¹, Edgar Omar Rueda Puente^{2*}, Onécimo Grimaldo Juárez¹, Daniel González Mendoza¹, Lourdes Cervantes Díaz¹ and Alejandro García López¹

¹Instituto de ciencias agrícolas, Universidad autónoma de Baja California. Carretera delta S/n Ejido Nuevo León Mexicali Baja California México

²Universidad de Sonora, Departamento de Agricultura y Ganadería. Carretera a Bahía de Kino, km. 21. Hermosillo, Sonora. México

*Address for Correspondence: Edgar Omar Rueda Puente, Instituto de ciencias agrícolas, Universidad autónoma de Baja California. Carretera delta S/n Ejido Nuevo León Mexicali Baja California México Cp. 21705, Tel: 6865230079; Email: erueda04@santana.uson.mx

Submitted: 22 December 2017

Approved: 02 January 2018

Published: 03 January 2018

Copyright: © 2018 Alzate JB, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Keywords: Stress; Abiotic; Propagation; Physiological-biochemical responses

Abstract

Vegetable production by grafting is a technique which it has made possible to resume agricultural soils which previously could not be produced due to stress generated by various abiotic factors, like a lack of water, stress by high or low temperatures, and or heavy metal contamination, among them. It has been documented and defined a number of graftings which they are tolerant to different factors; however, when it comes to auscultating information related to understand the molecular responses and observe what are the biochemical changes and physiological responses of grafted plants, it is dispersed. The current paper attempts to provide basic information documented on technique, addressing the molecular, biochemical and physiological responses, and thus get a clear perspective on the use of grafts, making this practice be used with most frequently by all its advantages.

Introduction

The first records of grafts showed that it was a long time in fruit trees [1]; During the 20s, the technique was used successfully in some vegetables [2], but it was not until the 21st century, the interest in its use for the production of large-scale crops [3]. The concern for the environment led to the search for alternatives in the integral production of crops, and one of them was the use of grafts. Currently there are some seed companies that have dedicated themselves to obtaining both improved crops and grafts. However, even though the advantage of the use of grafts is well known, several factors must be considered in order for the grafting technique to have the expected results. On the other hand, at present the tendency of the population is to consume more horticultural products, making the agricultural sector can not meet this demand without the use of large quantities of agrochemicals [4]. Along with this, the planting of products out of season and the use of land infested by phytopathogenic microorganisms generate a stress for the plants making frequent an imbalance in the normal development of the crops making necessary the use of agrochemicals and becoming an uncontrolled cycle that it damages the environment and generates several physiological disorders in plants and deterioration of the quality of the products consumed. For all this, is that the use of grafts has become an alternative to maintain the demand for horticultural products in many parts of the world.

Purpose of the grafts in vegetables

The main objective of the technique is the use of materials tolerant to diseases such as grafts, these materials make possible the production in soils where it would not normally be possible to normally. In some countries it is a widely used technique, although in the Occident it is a procedure recently introduced to the agricultural area. [5]. However, other important factors such as lack of water or the use of crops that tolerate different types of stress [6,7], make it necessary to search for alternatives such as grafts [8,9].

The grafting technique is usually done by hand, on the other hand, there is currently specialized equipment that is responsible for performing the grafts automatically. There are several types of grafts that, depending on the varieties to be grafted, the type of acclimatization chamber with which it is counted and the purposes for which the plant is wanted; the type of graft must be chosen. In figure 1, different variations of the technique for Solanaceae grafting are shown. Different technologies and procedures have been documented for other types of vegetables, which are described in the works of and [3,10].

This procedure has been used with great success for the control of diseases caused by fungi such as *Fusarium oxysporum*, *Sclerotium rolfsii*, *Verticillium*; wilting caused by bacteria such as *Ralstonia solanacearum* and some nematodes of the genus *Meloidogyne* spp. Among others [11-14]. However, the success of the technique for the management of diseases depends on a large number of factors; from the individual tolerance of grafts, the combination of variety- grafts and some other elements such as the genetic variability of plants [15,16].

Although these interactions and disease tolerance mechanisms are not yet fully understood, it is believed that the main tolerance is provided by the natural resistance of the grafts used. However, in studies where double-rooted grafts have been carried out, a partial tolerance of the grafted varieties is shown, suggesting that in the interaction between variety-grafts, associated substances are transported in defense against diseases such as *Fusarium*, said substances are synthesized in the root of the rootstock and transported via xylem to the variety [3,17]. Another of the main advantages of the use of grafts is the improvement in crop yields; Khanh et al. [18],




Figure 1: Grafting methods performed in Solanaceae; a) Double root approach method; b) tube insertion method; c) method of attachment; d) barb method; e) side pin insertion method. Bletsos and Olympios [61].

observed that tomatoes grafted on two hybrids as grafts had greater vigor than non-grafted varieties, in the same way they observed that the grafted plants had a higher yield ranging from 11 to 38% compared to non-grafted treatments. Similar results, but in watermelon, were obtained by Alan et al. [19], who observed a significant increase in the yield of grafted watermelon cultivated in two different environments, determine that under their study conditions the grafted watermelon not only increases agronomic parameters, but in the same way the technique does not affect the final quality of the watermelon fruit; this affirmation differs from that of Mohamed et al. [20], who also observed an increase in the yields of grafted watermelon. However, indicate that the fruit quality, especially the lycopene content in fruit, is modified under the conditions in which it was developed. The technique has also been used for other purposes such as improving the quality of grafted plants, greater absorption and intake of water and nutrients, tolerance to certain types of abiotic stress, tolerance to contamination by heavy metals, among others [21-24].

However, even knowing the advantages of its use, the main disadvantage of the technique of grafting, could be considered the economic factor, the use of tweezers or bands for gluing, knives, disinfectants, gloves, additional substrate, conditioning an acclimatization chamber, seed of the grafts (commercial hybrids), labor and other indirect factors cause the production costs of the grafted plant to rise [5], which in some cases could result in a possible disadvantage compared to the traditional production. Other factors to be considered as disadvantages may be the use of skilled labor, the lags in the production times, the initial delay caused by the grafting time and finally the limitation of the use of certain grafts.

Physiological, biochemical and molecular responses of grafted plants

The physiological responses of the grafted plants are given in the same way as in a normal plant, that is, they depend on factors such as the type of stress, duration of the same and intensity or progress of this, likewise it also depends on the genotype of the varieties, the state of development of the plants and their interaction with the environment [25,26].

Changes in the morphology of the grafted plant

Undoubtedly one of the most important aspects in grafted plants are the morphological modifications resulting from the grafts-variety interaction. Recently it was observed that grafted plants of beans had similar characteristics to non-grafted ones, however, factors such as plant height, number of leaves and flowers increased in plants grafted 28 days after the graft (DDI) Bernal-Alzate et al. [27], these modifications can be mediated by the production of growth regulators from the root transported via xylem to the aerial part of the plant [28,29]. On the other hand Kudo and Harada [30], observed how the morphology of the leaves of potato plants were modified due to the transport of RNA from the tomato rootstock towards the aerial part, determining that the amount of genetic material transported was enough to modify size, shape and quantity of trichomes in the leaves of grafted plants, these and other types of morphological changes have been documented by other researchers [31-33].

Stomatal conductance and CO₂ assimilation

Along with cell growth, photosynthesis is a primary process affected by lack of water and salinity, the effects on the plant range from a decrease in the assimilation of CO₂ caused by poor transport through stomata and mesophyll, to alterations in photosynthetic metabolism or even can lead to side effects causing oxidative stress in the cells [34-36], in a study conducted by Yang et al. [37], in plants of *Lagenaria siceraria* autografted and grafted on a rootstock tolerant to high salinity, it was observed that when increasing stress due to high salinity the stomatal conductivity (Gs) and the intracellular CO₂ concentration (Ci) decreased in both. However, those grafted on the

tolerant rootstock had higher Gs and Ci, indicating that probably stomatal closure was initiated by a signaling from the tolerant rootstock; These results are similar to those obtained by Rouphael et al. [38], who indicate that in both melon and grafted cucumbers there is a decrease in the photosynthetic activity of the plants. However, this decrease is up to 50% higher in plants that are not grafted and are under stress conditions due to high concentrations of NaCl; the same authors [38], found an inverse linear correlation of the photosynthetic activity with respect to the concentration of Na⁺ and Cl⁻ in leaves, attributing the excess of these to a disorder in the photosynthetic apparatus; similar studies witnessed the same photosynthetic activities [39,40]. Under normal conditions (without stress), grafted plants behave in a similar way to previous studies; in the case of Amaro et al. [41], observed that the grafting of the cucumber plants increased the stomatal conductance and photosynthetic capacity of the plant compared to those that had not been grafted; Similar results were reported by Liu et al. [42], attributing this increase in photosynthetic capacity to an increase in the amount of chlorophyll a and b in plants; additionally they reported the increase in carbohydrate accumulation in fruit, due to the activity of two key enzymes for this process in melon plants; Similar results were presented by QI et al. [43] and González et al. [44], in melon and citrus respectively. Likewise, Qinghai Gao et al. [45], Suggest that grafted plants have the ability to better use Cl under stress conditions, as well as presenting a better photosynthetic efficiency compared to those plants that were not grafted, this higher photosynthetic efficiency may be due to the fact that high levels of Cl decrease the transpiration of the leaves, improving the efficiency of water use in photosynthesis [46].

Enzymatic activity in grafts

When plants are under some kind of stress or when at some phenological stage of the crop occurs, one of the main indicators of these changes is the activity of certain enzymes. In grafts studies have been conducted to determine the activity of these enzymes. Some researchers have concentrated on observing that physiological and / or biochemical processes occur in different stages of the graft, it has been found that during the process of sticking the graft in incompatible plants there is an accumulation of ERO'S (Reactive oxygen species) [47-49], these in turn degrade the RUBISCO and therefore there is a decrease in the photosynthetic capacity of the plant specifically in carbon fixation [50]. Inefficiency in photosynthetic capacity leads to an incompatible plant death. Liao et al. [51], reported that the grafted promoted the increase in the concentration of key enzymes for RUBISCO photosynthesis (Ribulosa 1,5 biphosphate Carboxylase-Oxygenase) and RCA (RUBISCO activasa), these enzymes play an important role that directly affect the photosynthetic potential of the plants and therefore disrupt their yield potential [52]. Different types of response occur in grafted plants such as the accumulation of ROS, production of secondary metabolites directly or indirectly involved with the elimination of pathogens, synthesis of nitric oxide, and hypersensitive responses [53]. In the recent decade, studies have been carried out showing an increase in enzymatic activity and other biochemical compounds of grafted plants against various types of stress, Table 1 summarizes these studies.

Table 1: Enzymatic activity and other biochemical compounds of grafted plants against various types of stress.

Enzymes	Grafted plant	Condition of stress	Behavior of the enzyme	Reference
Catalase (CAT), Peroxidase (POD), Superoxide dismutase (SOD).	Tobacco	potato virus YNTN	Increase in POD and SOD, Decrease in CAT	[62]
Subunit 1 and 2 of the CAT isoenzyme	Tomato	<i>Fusarium</i> spp.	Presence of Subunits	[63]
CAT, SOD	Tomato	Water stress	SOD and CAT increase	[64]
Phenylalanine ammonium lyase (PAL)	Chili	<i>Fusarium</i> spp.	Increase at the beginning of the infection, stabilize or decrease days later.	[65]
Phenylalanine ammonium lyase (PAL)	Eggplant	<i>Verticillium</i> spp.	Increase PAL	[66]

Molecular responses of grafted plants

All physiological and / or biochemical responses are initialized by an expression of a specific gene. Work has been done on different types of grafts subjected to different types of stress to determine which specific genes are expressed and what is the biochemical or physiological response of these in grafted plants. It has been amply demonstrated that genetic exchange occurs in plants grafted differently, Cr  te et al. [54], establish that one of the factors for the exchange of silencing signals depends partially on the type of graft used, in their work they describe how using three different techniques of grafting only the technique of tube grafting (where a part of the variety wedge-shaped inserted in a ring cut on the rootstock), resulted in grafts systematically silenced by a sign of the rootstock; in 1997 Palauqui et al. [55], propose that the systemic silencing of genes in grafts occurs even when the expression of the silenced gene is not given in both varieties (rootstock and variety), these results do not agree with those reported by Shaharuddin et al. [56], who demonstrate that in tomato the exchange of messenger RNA (mRNA) between grafts-variety occurs only when the expression of a silenced gene is over expressed in both varieties before grafting. Wang et al. [57], found that the criollo varieties grafted tomato on a transgenic (which expresses the gene repressor MhGAI1 in the route of synthesis of gibberellins), accumulated transcripts of the gene MhGAI; these plants presented dwarfism, however, produced a greater amount of solids soluble in fruit, sugars and organic acids compared to those that were not grafted suggesting that the signaling for the synthesis of gibberellins, affected the quality of the fruit of the grafted plants. On the other hand, Ntatsi et al. [58], observed the tendency of accumulation of abscisic acid (ABA) in tomato plants was higher in grafted plants compared to non-grafted plants, regardless of the rootstock used in their study, attributing that the gene reviewed (LeNCED1), did not participate in the synthesis of ABA.

In a different work Jim  nez et al. [59], observed that the peach rootstock tolerant to salinity when subjected to this stress on expressed the specific gene P5SC at the same time that there was an increase of sorbitol, raffinose and proline determining that this gene could be used as a response marker to water stress in peach plants. One of the most recent studies on vegetables to date is that of Miao et al. [60], who based on the fact of the genetic transfer of the rootstock towards the variety, tested if there was a gene silencing related to the infection of the cucumber mosaic virus (CMV Cucumber mosaic virus); they tested a transgenic rootstock resistant to the virus and a susceptible variety, finding that the gene silencing provided by the resistance to this virus was transmitted by the rootstock towards the variety; the same work concluded, that these plants may remain undetectable for gene markers of transgenic resistance and, may provide new approaches to evade concerns with biosecurity over genetically modified organisms (GMOs).

Conclusions

The use of grafts is a relatively new technique in the agricultural area, the increase of its use is due in large part to its multiple advantages outweigh the economic factor that entails. That is why currently it is necessary to focus part of the research in this area to the search of Creole grafts that reduce the costs of the technique; This search should be conducted in finding materials not only that lead to increase the productive potential of the plants, but also, the grafts obtained must be tolerant to some type of stress. It is undeniable that the use of the technique represents an alternative for the production of vegetables in a way more friendly to the environment, so its use could be increased in the coming years.

On the other hand, at the present time the increasing number of researches to enlarge of knowledge in the understanding of the physiological, biochemical and molecular changes that occur in grafted plants, continues to rise and although more and more is understood about the behavior of grafts under different types of conditions; sometimes the results of these investigations may not coincide and sometimes not be conclusive.

Acknowledgments

To the Universidad Autónoma de Baja California (UABC), for the support granted through the 17th Internal Call for Research Projects and doctoral fellowship provided by the National Council of Science and Technology.

References

1. Lockard RG. Effect of Apple Rootstocks and Length and Type of Interstock on Leaf Nutrient Levels. *J Hortic Sci.* 1976; 51: 289-296. [Ref.: https://goo.gl/kuE7q5](https://goo.gl/kuE7q5)
2. Tateishi K. Grafting watermelon onto pumpkin. *J Japanese Horticulture (Nihon-Engei Zasshi).* 1927; 39: 5-8. [Ref.: https://goo.gl/jKnxRD](https://goo.gl/jKnxRD)
3. Lee JM, Kubota C, Tsao SJ, Bie Z, Hoyos-Echeverria P, et al. Current status of vegetable grafting: Diffusion, grafting techniques, automation. *Scientia Horticulturae.* 2010; 127: 93-105. [Ref.: https://goo.gl/P93mPv](https://goo.gl/P93mPv)
4. OCDE/FAO (2013), OCDE-FAO Perspectivas Agrícolas 2013-2022, Texcoco, Estado de México, Universidad Autónoma Chapingo.
5. Louws FJ, Rivard CL, Kubota C. Grafting fruiting vegetables to manage soilborne pathogens, foliar pathogens, arthropods and weeds. *Sci Hortic.* 2010; 127: 127-146. [Ref.: https://goo.gl/DvV4gA](https://goo.gl/DvV4gA)
6. Luna-Flores W, Estrada-Medina H, Jiménez-Osornio JJM, Pinzón-López LL. Efecto del estrés hídrico sobre el crecimiento y eficiencia del uso del agua en plántulas de tres especies arbóreas caducifolias. *Terra Latinoamericana.* 2012; 30: 343-353. [Ref.: https://goo.gl/QeE3su](https://goo.gl/QeE3su)
7. Moreno L. Respuesta de las plantas al estrés por déficit hídrico. Una revisión. *Agronomía colombiana.* 2009; 27: 179-191. [Ref.: https://goo.gl/F1Zt4t](https://goo.gl/F1Zt4t)
8. Penella C, Landi M, Guidi L, Nebauer S, Pellegrini E, et al. Salt-tolerant rootstock increases yield of pepper under salinity through maintenance of photosynthetic performance and sinks strength. *J plant Physiology.* 2016; 193: 1-11. [Ref.: https://goo.gl/MEhCMF](https://goo.gl/MEhCMF)
9. Sánchez-Rodríguez E, Rubio-Wilhelmi M, Blasco B, Leyva R, Romero L, et al. Antioxidant response resides in the shoot in reciprocal grafts of drought-tolerant and drought-sensitive cultivars in tomato under water stress. *Plant Science.* 2012; 189: 89-96. [Ref.: https://goo.gl/AViWS9](https://goo.gl/AViWS9)
10. Kubota C, Mc Clure M, Kokalis-Burelle N, Bausher M, Roskopf E. Vegetable grafting: History, use and current technology status in North America. *HortScience.* 2008; 6: 1664-1669. [Ref.: https://goo.gl/ytevSe](https://goo.gl/ytevSe)
11. Bletsos F, Thanassouloupoulos C, Roupakias D. Effect of grafting on growth, yield, and verticillium wilt of eggplant. *HortScience.* 2003; 2: 183-186. [Ref.: https://goo.gl/5PMJIF](https://goo.gl/5PMJIF)
12. Rivard C, O'Connell S, Peet M, Louws F. Grafting tomato with interspecific Rootstock to manage diseases caused by *Sclerotium rolfsii* and southern root-knot nematode. *Plant disease.* 2010; 8: 1015-1021. [Ref.: https://goo.gl/JyVCfw](https://goo.gl/JyVCfw)
13. Rivard C, O'Connell S, Peet M, Welker R, Louws F. Grafting tomato to manage bacterial wilt caused by *Ralstonia solanacearum* in the southeastern United States. *Plant disease.* 2012; 7: 973-978.
14. Keinath A, Haseell R. Control of fusarium wilt of watermelon by grafting onto bottlegourd of interspecific hybrid squash despite colonization of grafts by *Fusarium*. *Plants disease.* 2014; 2: 255-266. [Ref.: https://goo.gl/TVJYcX](https://goo.gl/TVJYcX)
15. Kleinhenz MD. Major Factors in Preparing Grafted Vegetable Plants Successfully. The Ohio State Univ., Ohio Agricultural Res. Dev. Ctr. 2011.
16. Ozores-Hampton M. Healing chamber for grafted vegetables seedlings in Florida. University of Florida IFAS. 2013.
17. Dawson R. Accumulation of nicotine in reciprocal grafts of tomato and tobacco. *American Journal of botany.* 1942; 29: 66-71. [Ref.: https://goo.gl/byZMhm](https://goo.gl/byZMhm)
18. Khah E, Kakava E, Mavromatis A, Chachalis D, Goulas C. Effect of grafting on growth and yield of tomato (*Lycopersicon esculentum* Mill.) in greenhouse and open field. *Journal of Applied Horticulture.* 2006; 8: 3-7. [Ref.: https://goo.gl/tTkW3L](https://goo.gl/tTkW3L)
19. Alan Ö, Özdemir N, Günen Y. Effect of grafting on watermelon plant growth, yield and quality. *Journal of Agronomy.* 2007; 2: 362-365. [Ref.: https://goo.gl/bxbwcX](https://goo.gl/bxbwcX)

20. Mohamed F, El-Hamed K, Elwhan M, Hussien M. Impact of grafting on watermelon growth, fruit yield and quality. *Vegetable Research Bulletin*. 2012; 76: 99-118. [Ref.:](https://goo.gl/of3LLz) <https://goo.gl/of3LLz>
21. Savvas D, Colla G, Roupael Y, Schwarz D. Amelioration of heavy metal and nutrient stress in fruit vegetables by grafting. *Scientia Horticulturae*. 2010; 2: 156-161. [Ref.:](https://goo.gl/q3PcNM) <https://goo.gl/q3PcNM>
22. Di Gioia F, Signore A. Grafting improves tomato salinity tolerance through sodium partitioning within the shoot. *HortScience*. 2013; 7: 855-862. [Ref.:](https://goo.gl/dmX6YM) <https://goo.gl/dmX6YM>
23. Sánchez-Rodríguez E, Romero L, Ruiz JM. Role of grafting in resistance to water stress in tomato plants: ammonia production and assimilation. *J Plant Growth Regul*. 2013; 32: 831-842. [Ref.:](https://goo.gl/fUHyJz) <https://goo.gl/fUHyJz>
24. Savvas D, Ntatsi G, Barouchas P. Impact of grafting and rootstock genotype on cation uptake by cucumber (*Cucumis sativus* L.) exposed to Cd or Ni stress. *Scientia Horticulturae*. 2013; 149: 86-96. [Ref.:](https://goo.gl/7XBwkZ) <https://goo.gl/7XBwkZ>
25. Chaves M, Maroco J, Pereira. Understanding plant responses to drought -from genes to the whole plant. *Funct Plant Biol*. 2003; 30: 239-264. [Ref.:](https://goo.gl/Umca4W) <https://goo.gl/Umca4W>
26. Blum A. Drought resistance, water use-efficiency, and yield potential -are they compatible, dissonant or mutual exclusive? *Austr J Agric Res*. 2005; 56: 1159-1168. [Ref.:](https://goo.gl/zkxDLQ) <https://goo.gl/zkxDLQ>
27. Bernal-Alzate J, Grimaldo-Juarez O, González-Mendoza D, Cervantes-Díaz L, Rueda-Puente E, et al. El injerto como alternativa para mejorar el rendimiento en la producción de frijol ejotero (*Phaseolus vulgaris* L.). *IDESIA*. 2016; 2: 43-46. [Ref.:](https://goo.gl/pZWYko) <https://goo.gl/pZWYko>
28. Proebsting W, Hedden P, Lewis M, Croker S, Proebsting L. Gibberellin concentration and transport in genetic lines of pea. *Plant Physiol*. 1992; 100: 1354-1360. [Ref.:](https://goo.gl/TSzCDg) <https://goo.gl/TSzCDg>
29. Bulley S, Wilson F, Hedden P, Phillips A, Croker S, et al. Modification of gibberellin biosynthesis in the grafted apple scion allows control of tree height independent of the rootstock. *Plant Biotechnology Journal*. 2005; 3: 215-223. [Ref.:](https://goo.gl/w2HBW6) <https://goo.gl/w2HBW6>
30. Kudo H, Harada T. A graft-transmissible RNA from Tomato Rootstock changes leaf morphology of potato scion. *Hortscience*. 2007; 2: 225-226. [Ref.:](https://goo.gl/in7stW) <https://goo.gl/in7stW>
31. Ohata Y. Graft-transformation, the mechanism for graft-induced genetic changes in higher plants. *Euphytica*. 1991; 55: 91-99. [Ref.:](https://goo.gl/Bi1J6J) <https://goo.gl/Bi1J6J>
32. Taller J, Yagishita N, Hirata Y. Graft-induced variants as a source of novel characteristics in the breeding pepper (*Capsicum annuum* L.). *Euphytica*. 1999; 108: 73-78. [Ref.:](https://goo.gl/X2P66N) <https://goo.gl/X2P66N>
33. Hooijdonk B, Woolley D, Warrington I, Tustin S. Rootstocks modify scion architecture, endogenous hormones and root growth of newly grafted 'royal gala' apple trees. *J. Amer. Soc. Hort. Sci*. 2011; 136: 93-102. [Ref.:](https://goo.gl/Kobncn) <https://goo.gl/Kobncn>
34. Sandalio L, Dalurzo H, Gimez M, Romero-Puertas M, Rio L. Cadmium-induced changes in growth and oxidative metabolism of pea plants, *J. Exp. Bot*. 2001; 52: 1297-1303. [Ref.:](https://goo.gl/ETwErt) <https://goo.gl/ETwErt>
35. Saied A, Keutgen N, Noga G. Effects of NaCl stress on leaf growth, photosynthesis and ionic contents of strawberry cvs 'Elsanta' and 'Korona'. In: pardossi, A., Serra, G., F. (Eds.). *International symposium on managing greenhouse crops in saline environment*, International society of Horticultural Science. Pisa: 2003; 67-73.
36. Chaves M, Flexas J, Pinheiro C. Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Annals of botany*. 2009; 103: 551-560. [Ref.:](https://goo.gl/MHXnwi) <https://goo.gl/MHXnwi>
37. Yuang Y, Yu L, Wang L, Guo S. Bottle gourd grafts-grafting promotes photosynthesis by regulating the stomata and non-stomata performances in leaves of watermelon seedlings under NaCl stress. *Journal of plant Physiology*. 2015; 187: 50-58. [Ref.:](https://goo.gl/6KFERD) <https://goo.gl/6KFERD>
38. Roupael Y, Cardarelli M, Rea E, Colla G. Improving melon and cucumber photosynthetic activity, mineral composition, and growth performance under salinity stress by grafting onto *Cucurbita* hybrid grafts. *Photosynthetica*. 2009; 50: 180-188. [Ref.:](https://goo.gl/afym6X) <https://goo.gl/afym6X>
39. Aghaleh M, Niknam V, Ebrahimzadeh H, Razavi K. Salt stress effects on growth, pigments, proteins and lipid peroxidation in *Salicornia persica* and *S. europaea* *Biol Plant*. 2009; 53: 243-248. [Ref.:](https://goo.gl/p3KztJ) <https://goo.gl/p3KztJ>
40. Liu Z, Bie Z, Huang Y, Zhen A, Lei B, et al. Grafting onto *Cucurbita moschata* rootstock alleviates salt stress in cucumber plants by delaying photoinhibition. *Photosynthetica*. 2012; 50: 152-160. [Ref.:](https://goo.gl/hxJ8WV) <https://goo.gl/hxJ8WV>



41. Amaro A, Macedo A, Pereira A, Goto R, Ono E, et al. The use of grafting to improve the net photosynthesis of cucumber. *Theor exp. Plant Physiol.* 2014; 26: 241-249. [Ref.: https://goo.gl/3jQsrU](https://goo.gl/3jQsrU)
42. Liu Y, Qi Y, Bai M, Qi F, Xu Q, et al. Grafting Helps Improve Photosynthesis and Carbohydrate Metabolism in Leaves of Muskmelon. *Int J Biol Sci.* 2011; 7: 1161-1170. [Ref.: https://goo.gl/B7eyp9](https://goo.gl/B7eyp9)
43. Qi Y, Li L, Liu F, Li D. Effects of grafting on photosynthesis characteristics, yield and sugar content in melon. *J Shenyang Agr Univ.* 2006; 37: 155-158. [Ref.: https://goo.gl/zSShUJ](https://goo.gl/zSShUJ)
44. Gonzalez C, Llosa J, Quijano A, Forner A. Rootstock effects on leaf Photosynthesis in "Navelina" Trees grown in calcareous soil. *HortScience.* 2009; 44: 280-283. [Ref.: https://goo.gl/Rmt6vh](https://goo.gl/Rmt6vh)
45. Qinghai G, Wu Y, Jia S, Huang S, Lu X. Effect of rootstock on the growth, photosynthetic capacity And osmotic adjustment of eggplant seedlings under Chilling stress and recovery. *Pak. J. Bot.* 2016; 48: 461-467. [Ref.: https://goo.gl/83WUxQ](https://goo.gl/83WUxQ)
46. Jianlin W Y, Guirui F, Quanxiao J, Defeng Q, Hua W, et al. Responses of water use efficiency of 9 plant species to light and CO₂ and their modeling. *Acta Ecol.* 2008; 28: 525-533. [Ref.: https://goo.gl/9n9C6y](https://goo.gl/9n9C6y)
47. Aloni B, Karni L, Deventurero G, Levin Z, Cohen R, et al. Physiological and biochemical changes at the grafts-scion interface in graft combinations between Cucurbita grafts and a melon scion. *J. Hortic. Sci. Biotechnol.* [Ref.: 2008; 83: 777-783. Ref.: https://goo.gl/wJypQy](https://goo.gl/wJypQy)
48. Irisarri P, Binczycki P, Errea P, Martens H J, Pina A. Oxidative stress associated with rootstockescion interactions in pear/quince combinations during early stages of graft development. *J. Plant Physiol.* 2015; 176: 25-35. [Ref.: https://goo.gl/TsNPHy](https://goo.gl/TsNPHy)
49. Xu Q, GHuo S, Li L, An Y, Shu S, et al. Proteomics analysis of compatibility and incompatibility in grafted cucumber seedlings. *Plants physiology and biochemistry.* 2016; 105: 21-28. [Ref.: https://goo.gl/2fQgWD](https://goo.gl/2fQgWD)
50. Desimone M, Henke A, Wagner E. Oxidative stress induces partial degradation of the large subunit of ribulose-1, 5-bisphosphate carboxylase/oxygenase in isolated chloroplasts of barley. *Plant Physiol.* 1996; 111: 789-796. [Ref.: https://goo.gl/FFLcfD](https://goo.gl/FFLcfD)
51. Liao L, Cao S, Rong Y, Wang Z. Effects of grafting on key photosynthetic enzymes and gene expression in the citrus cultivar Huangguogan. *Genetics and molecular research.* 2016; 15: 1-10. [Ref.: https://goo.gl/KrVQZy](https://goo.gl/KrVQZy)
52. Morinaga K, Ikeda F. The effects of several grafts on photosynthesis; distribution of photosynthetic product, and growth of young satsuma mandarin trees. *J. Japan. Soc. Hort. Sci.* 1990; 59: 29-34. [Ref.: https://goo.gl/4XuGDj](https://goo.gl/4XuGDj)
53. Buchanan B, W Gruissem, Jones R. *Biochemistry and molecular biology of plants.* American Society of Plant Biologists, John Wiley & Sons, Inc. Somerset NJ. 2000. [Ref.: https://goo.gl/T5fodG](https://goo.gl/T5fodG)
54. Cr  t   P, Leuenberger S, V A Iglesias, V Suarez, H Schob, et al. Graft transmission of induced and spontaneous post-transcriptional silencing of chitinase genes. *Plant J.* 2001; 28: 493-501. [Ref.: https://goo.gl/3A6J2c](https://goo.gl/3A6J2c)
55. Palauqui J C, Elmayan T, Pollien J M, Vaucheret H. Systemic acquired silencing: transgene-specific posttranscriptional silencing is transmitted by grafting from silenced stocks to non-silenced scions. *EMBO J.* 1997; 15: 4738-4745. [Ref.: https://goo.gl/wqbsJX](https://goo.gl/wqbsJX)
56. Shaharuddin N, Han Y, Li H, Grierson D. The mechanism of graft transmission of sense and antisense gene silencing in tomato plants. *FEBS letters.* 2006; 580: 6579-6586. [Ref.: https://goo.gl/vhkY9f](https://goo.gl/vhkY9f)
57. Wang S, Liu Z, Sun C, Shi Q, Yao Y, et al. Functional characterization of the apple *MhGAI1* gene through ectopic expression and grafting experiments in tomatoes. *Journal of plant physiology.* 2012; 169: 303-310. [Ref.: https://goo.gl/KQ9GDb](https://goo.gl/KQ9GDb)
58. Ntatsi G, Savvas D, Huntenturg K, Druege U, Hinch D, et al. A study on ABA involvement in the response of tomato suboptimal temperatura using reciprocal grafts with *notabilis*, a null mutant in the ABA-biosynthesis gene *LeNCED*. *Enviromental and Experimental Botany.* 2014; 97: 11-21.
59. Jim  nez S, Dridi J, Guti  rrez D, Moret D, Irigoyen J, et al. Physiological, biochemical and molecular responses of four *prunus* grafts submitted to drought stress. *Tree physiology.* 2013; 33: 1061-1075. [Ref.: https://goo.gl/tWefsZ](https://goo.gl/tWefsZ)
60. Miao B, Wen-ting C, Bing-yan X, Guo-shun Y. A novel strategy to enhance resistance to *Cucumber mosaic virus* in tomato by grafting to transgenic grafts. *Journal of integrative agriculture.* 2016; 15: 2040-2048. [Ref.: https://goo.gl/JdXz4H](https://goo.gl/JdXz4H)



61. Bletsos F, Olympios C. Grafts and grafting of tomatoes, peppers and eggplants for soil-borne disease resistance, improved yield and quality. *The European journal of plant science and biotechnology*. 2008; 2: 62-73.
62. Spoustová P, Hýsková V, Müller K, Schnablová R, Ryslavá H, et al. Tobacco susceptibility to Potato virus Y^{NTN} infection is affected by grafting and endogenous cytokinin content. *Plant Science*. 2015; 235: 25-36. [Ref.: https://goo.gl/omxqFf](https://goo.gl/omxqFf)
63. Vitale A, Rocco M, Arena S, Giuffrida F, Cassanitu C, et al. Tomato susceptibility to Fusarium crown and root rot: Effect of grafting combination and proteomic analysis of tolerance expression in the rootstock. *Plant Physiology and biochemistry*. 2014; 83: 207- 216. [Ref.: https://goo.gl/bsqrtA](https://goo.gl/bsqrtA)
64. Sánchez-Rodríguez, E Ruiz, J Ferreres, F Moreno, D. Phenolic profiles of cherry tomatoes as influenced by hydric stress and rootstock technique. *Food chemistry*. 2012; 134: 775-782. [Ref.: https://goo.gl/Smx5up](https://goo.gl/Smx5up)
65. Jiang F, Y X Liu, W Liu, N Zheng, H T Wang, et al. Relationship between root rot resistance and phenylalanine metabolism in graft capsicum. *China Veg*. 2010; 8: 46-52. [Ref.: https://goo.gl/Bxw8Lt](https://goo.gl/Bxw8Lt)
66. Zhou B, Gao Y, Lin G, Fu Y. Relationship between disease resistance and electrolytic leakage, proline content and PAL activity in grafted eggplant (in Chinese). *Acta Hort Sinica* 1998; 25: 300-302. [Ref.: https://goo.gl/6m8eCs](https://goo.gl/6m8eCs)
67. Edelstein M, Cohen R, Burger Y, Shriber S, Pivonia S, et al. Integrated management of sudden wilt of melons, caused by *Monosporascus cannonballus*, using grafting and reduced rate of methyl bromide. *Plant Dis*. 1999; 83: 1142-1145. [Ref.: https://goo.gl/XPx9Rn](https://goo.gl/XPx9Rn)
68. Franks. P, Casson S. Connecting stomatal development and physiology. *New Phytol*. 2014; 201: 1079-1082. [Ref.: https://goo.gl/Btu4ir](https://goo.gl/Btu4ir)
69. Jones H. *Plants and microclimate: a Quantitative Approach to Environmental*. Plant physiology, 3ra Edicion. Cambridge University Press London. 2014. [Ref.: https://goo.gl/E5mZNs](https://goo.gl/E5mZNs)
70. Rivard C, Sydorovych O, O'Connell S, Peet M, Louws F. An economic analysis of two grafted tomato transplant production systems in the United States. *Horttechnology*. 2010; 4: 794-803. [Ref.: https://goo.gl/vdwYlf](https://goo.gl/vdwYlf)